

- Introduction to Krylov subspace techniques
- FOM, GMRES, practical details.
- Symmetric case: Conjugate gradient
- See Chapter 6 of text for details.

Motivation

- Common feature of one-dimensional projection techniques:

$$x_{new} = x + \alpha d$$

where d = a certain direction.

- α is defined to optimize a certain function.
- Equivalently: determine α by an orthogonality constraint

Example


In MR:
 $x(\alpha) = x + \alpha d$, with $d = b - Ax$.
 $\min_{\alpha} \|b - Ax(\alpha)\|_2$ reached iff $b - Ax(\alpha) \perp r$

- One-dimensional projection methods are greedy methods. They are 'short-sighted'.

Example:

Recall in Steepest Descent: New direction of search \tilde{r} is \perp to old direction of search r .

$$\begin{aligned} r &\leftarrow b - Ax, \\ \alpha &\leftarrow (r, r) / (Ar, r) \\ x &\leftarrow x + \alpha r \end{aligned}$$



Question: can we do better by combining successive iterates?

- Yes: Krylov subspace methods..

Krylov subspace methods: Introduction

- Consider MR (or steepest descent). At each iteration:

$$\begin{aligned} r_{k+1} &= b - A(x^{(k)} + \alpha_k r_k) \\ &= r_k - \alpha_k A r_k \\ &= (I - \alpha_k A) r_k \end{aligned}$$

- In the end:

$$r_{k+1} = (I - \alpha_k A)(I - \alpha_{k-1} A) \cdots (I - \alpha_0 A) r_0 = p_{k+1}(A) r_0$$

where $p_{k+1}(t)$ is a polynomial of degree $k + 1$ of the form

$$p_{k+1}(t) = 1 - tq_k(t)$$

- ☞ Show that: $x^{(k+1)} = x^{(0)} + q_k(A)r_0$, with $\deg(q_k) = k$

- Krylov subspace methods: iterations of this form that are 'optimal' [from m -dimensional projection methods]

Krylov subspace methods

Principle: Projection methods on Krylov subspaces:

$$K_m(A, v_1) = \text{span}\{v_1, Av_1, \dots, A^{m-1}v_1\}$$

- The most important class of iterative methods.
- Many variants exist depending on the subspace L .

Simple properties of K_m

► Notation: $\mu = \text{deg. of minimal polynomial of } v$. Then:

- $K_m = \{p(A)v \mid p = \text{polynomial of degree } \leq m - 1\}$
- $K_m = K_\mu$ for all $m \geq \mu$. Moreover, K_μ is invariant under A .
- $\dim(K_m) = m$ iff $\mu \geq m$.

13-5

Text: 6 – Krylov1

A little review: Gram-Schmidt process

Goal: given $X = [x_1, \dots, x_m]$ compute an orthonormal set $Q = [q_1, \dots, q_m]$ which spans the same subspace.

ALGORITHM : 1. Classical Gram-Schmidt

1. For $j = 1, \dots, m$ Do:
2. Compute $r_{ij} = (x_j, q_i)$ for $i = 1, \dots, j - 1$
3. Compute $\hat{q}_j = x_j - \sum_{i=1}^{j-1} r_{ij}q_i$
4. $r_{jj} = \|\hat{q}_j\|_2$ If $r_{jj} == 0$ exit
5. $q_j = \hat{q}_j / r_{jj}$
6. EndDo

13-6

Text: 6 – Krylov1

ALGORITHM : 2. Modified Gram-Schmidt

1. For $j = 1, \dots, m$ Do:
2. $\hat{q}_j := x_j$
3. For $i = 1, \dots, j - 1$ Do
4. $r_{ij} = (\hat{q}_j, q_i)$
5. $\hat{q}_j := \hat{q}_j - r_{ij}q_i$
6. EndDo
7. $r_{jj} = \|\hat{q}_j\|_2$. If $r_{jj} == 0$ exit
8. $q_j := \hat{q}_j / r_{jj}$
9. EndDo

13-7

Text: 6 – Krylov1

Let:

$X = [x_1, \dots, x_m]$ ($n \times m$ matrix)

$Q = [q_1, \dots, q_m]$ ($n \times m$ matrix)

$R = \{r_{ij}\}$ ($m \times m$ upper triangular matrix)

► At each step,

$$x_j = \sum_{i=1}^j r_{ij}q_i$$

Result:

$$X = QR$$

13-8

Text: 6 – Krylov1

$$\bar{H}_m^{(1)} = \begin{bmatrix} h_{11}^{(1)} & h_{12}^{(1)} & h_{13}^{(1)} & h_{14}^{(1)} & h_{15}^{(1)} \\ & h_{22}^{(1)} & h_{23}^{(1)} & h_{24}^{(1)} & h_{25}^{(1)} \\ & h_{32} & h_{33} & h_{34} & h_{35} \\ & & h_{43} & h_{44} & h_{45} \\ & & & h_{54} & h_{55} \\ & & & & h_{65} \end{bmatrix}, \bar{g}_1 = \begin{bmatrix} c_1\beta \\ -s_1\beta \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Repeat with Ω_2 ,
 \dots ,
 Ω_5 .
 Result:

$$\bar{H}_5^{(5)} = \begin{bmatrix} h_{11}^{(5)} & h_{12}^{(5)} & h_{13}^{(5)} & h_{14}^{(5)} & h_{15}^{(5)} \\ & h_{22}^{(5)} & h_{23}^{(5)} & h_{24}^{(5)} & h_{25}^{(5)} \\ & & h_{33}^{(5)} & h_{34}^{(5)} & h_{35}^{(5)} \\ & & & h_{44}^{(5)} & h_{45}^{(5)} \\ & & & & h_{55}^{(5)} \\ & & & & 0 \end{bmatrix}, \bar{g}_5 = \begin{bmatrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \cdot \\ \cdot \\ \gamma_6 \end{bmatrix}$$

13-17

Text: 6 – Krylov1

Define

$$Q_m = \Omega_m \Omega_{m-1} \dots \Omega_1$$

$$\bar{R}_m = \bar{H}_m^{(m)} = Q_m \bar{H}_m,$$

$$\bar{g}_m = Q_m(\beta e_1) = (\gamma_1, \dots, \gamma_{m+1})^T.$$

➤ Since Q_m is unitary,

$$\min \|\beta e_1 - \bar{H}_m y\|_2 = \min \|\bar{g}_m - \bar{R}_m y\|_2.$$

➤ Delete last row and solve resulting triangular system.

$$R_m y_m = g_m$$

13-18

Text: 6 – Krylov1

Proposition:

1. The rank of AV_m is equal to the rank of R_m . In particular, if $r_{mm} = 0$ then A must be singular.
2. The vector y_m that minimizes $\|\beta e_1 - \bar{H}_m y\|_2$ is given by

$$y_m = R_m^{-1} g_m.$$

3. The residual vector at step m satisfies

$$\begin{aligned} b - Ax_m &= V_{m+1} [\beta e_1 - \bar{H}_m y_m] \\ &= V_{m+1} Q_m^T (\gamma_{m+1} e_{m+1}) \end{aligned}$$

4. As a result, $\|b - Ax_m\|_2 = |\gamma_{m+1}|$.

13-19

Text: 6 – Krylov1