

ERROR AND SENSITIVITY ANALYSIS FOR SYSTEMS OF LINEAR EQUATIONS

- Conditioning of linear systems.
- Estimating errors for solutions of linear systems
- (Normwise) Backward error analysis
- Estimating condition numbers ..

5-1

Perturbation analysis for linear systems ($Ax = b$)

Question addressed by perturbation analysis: determine the variation of the solution x when the data, namely A and b , undergoes small variations. Problem is **Ill-conditioned** if small variations in data cause very large variation in the solution.

Setting:

► We perturb A into $A + E$ and b into $b + e_b$. Can we bound the resulting change (perturbation) to the solution?

Preparation: We begin with a lemma for a simple case

5-2

TB: 12; AB: 1.2.7; GvL 3.5 – PertA

5-2

Rigorous norm-based error bounds

LEMMA: If $\|E\| < 1$ then $I - E$ is nonsingular and

$$\|(I - E)^{-1}\| \leq \frac{1}{1 - \|E\|}$$

Proof is based on following 5 steps

- a) Show: If $\|E\| < 1$ then $I - E$ is nonsingular
- b) Show: $(I - E)(I + E + E^2 + \dots + E^k) = I - E^{k+1}$.
- c) From which we get:

$$(I - E)^{-1} = \sum_{i=0}^k E^i + (I - E)^{-1} E^{k+1} \rightarrow$$

5-3

d) $(I - E)^{-1} = \lim_{k \rightarrow \infty} \sum_{i=0}^k E^i$. We write this as

$$(I - E)^{-1} = \sum_{i=0}^{\infty} E^i$$

e) Finally:

$$\begin{aligned} \|(I - E)^{-1}\| &= \left\| \lim_{k \rightarrow \infty} \sum_{i=0}^k E^i \right\| = \lim_{k \rightarrow \infty} \left\| \sum_{i=0}^k E^i \right\| \\ &\leq \lim_{k \rightarrow \infty} \sum_{i=0}^k \|E^i\| \leq \lim_{k \rightarrow \infty} \sum_{i=0}^k \|E\|^i \\ &\leq \frac{1}{1 - \|E\|} \end{aligned}$$

5-4

TB: 12; AB: 1.2.7; GvL 3.5 – PertA

5-4

- Can generalize result:

LEMMA: If A is nonsingular and $\|A^{-1}\| \|E\| < 1$ then $A + E$ is non-singular and

$$\|(A + E)^{-1}\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1}\| \|E\|}$$

- Proof is based on relation $A + E = A(I + A^{-1}E)$ and use of previous lemma.
- Now we can prove the main theorem:

THEOREM 1: Assume that $(A + E)y = b + e_b$ and $Ax = b$ and that $\|A^{-1}\| \|E\| < 1$. Then $A + E$ is nonsingular and

$$\frac{\|x - y\|}{\|x\|} \leq \frac{\|A^{-1}\| \|A\|}{1 - \|A^{-1}\| \|E\|} \left(\frac{\|E\|}{\|A\|} + \frac{\|e_b\|}{\|b\|} \right)$$

Proof: From $(A + E)y = b + e_b$ and $Ax = b$ we get $(A + E)(y - x) = e_b - Ex$. Hence:

$$y - x = (A + E)^{-1}(e_b - Ex)$$

Taking norms $\rightarrow \|y - x\| \leq \|(A + E)^{-1}\| [\|e_b\| + \|E\| \|x\|]$
Dividing by $\|x\|$ and using result of lemma

$$\begin{aligned} \frac{\|y - x\|}{\|x\|} &\leq \|(A + E)^{-1}\| [\|e_b\|/\|x\| + \|E\|] \\ &\leq \frac{\|A^{-1}\|}{1 - \|A^{-1}\| \|E\|} [\|e_b\|/\|x\| + \|E\|] \\ &\leq \frac{\|A^{-1}\| \|A\|}{1 - \|A^{-1}\| \|E\|} \left[\frac{\|e_b\|}{\|A\| \|x\|} + \frac{\|E\|}{\|A\|} \right] \end{aligned}$$

Result follows by using inequality $\|A\| \|x\| \geq \|b\| \dots$ **QED**

The quantity $\kappa(A) = \|A\| \|A^{-1}\|$ is called the **condition number** of the linear system with respect to the norm $\|\cdot\|$. When using the p -norms we write:

$$\kappa_p(A) = \|A\|_p \|A^{-1}\|_p$$

- Note: $\kappa_2(A) = \sigma_{max}(A)/\sigma_{min}(A)$ = ratio of largest to smallest singular values of A . Allows to define $\kappa_2(A)$ when A is not square.
- Determinant *is not* a good indication of sensitivity
- Small eigenvalues *do not* always give a good indication of poor conditioning.

Example: Consider, for a large α , the $n \times n$ matrix

$$A = I + \alpha e_1 e_n^T$$

- Inverse of A is : $A^{-1} = I - \alpha e_1 e_n^T$ ➤ For the ∞ -norm we have

$$\|A\|_\infty = \|A^{-1}\|_\infty = 1 + |\alpha|$$

so that $\kappa_\infty(A) = (1 + |\alpha|)^2$.

- Can give a very large condition number for a large α – but all the eigenvalues of A are equal to one.

- ☞₁ Show that $\kappa(I) = 1$;
- ☞₂ Show that $\kappa(A) \geq 1$;
- ☞₃ Show that $\kappa(A) = \kappa(A^{-1})$
- ☞₄ Show that for $\alpha \neq 0$, we have $\kappa(\alpha A) = \kappa(A)$

Simplification when $e_b = 0$:	Simplification when $E = 0$:
$\frac{\ x - y\ }{\ x\ } \leq \frac{\ A^{-1}\ \ E\ }{1 - \ A^{-1}\ \ E\ }$	$\frac{\ x - y\ }{\ x\ } \leq \ A^{-1}\ \ A\ \frac{\ e_b\ }{\ b\ }$

➤ Slightly less general form: Assume that $\|E\|/\|A\| \leq \delta$ and $\|e_b\|/\|b\| \leq \delta$ and $\delta\kappa(A) < 1$ then

$$\frac{\|x - y\|}{\|x\|} \leq \frac{2\delta\kappa(A)}{1 - \delta\kappa(A)}$$

☞₅ Show the above result

Another common form:

THEOREM 2: Let $(A + \Delta A)y = b + \Delta b$ and $Ax = b$ where $\|\Delta A\| \leq \epsilon\|E\|$, $\|\Delta b\| \leq \epsilon\|e_b\|$, and assume that $\epsilon\|A^{-1}\|\|E\| < 1$. Then

$$\frac{\|x - y\|}{\|x\|} \leq \frac{\epsilon \|A^{-1}\| \|A\|}{1 - \epsilon\|A^{-1}\| \|E\|} \left(\frac{\|e_b\|}{\|b\|} + \frac{\|E\|}{\|A\|} \right)$$

➤ Results to be seen later are of this type.

Normwise backward error

➤ We solve $Ax = b$ and find an approximate solution y

Question: Find smallest perturbation to apply to A, b so that *exact* solution of perturbed system is y

Normwise backward error in just A or b

Suppose we model entire perturbation in RHS b .

- Let $r = b - Ay$ be the residual.
Then y satisfies $Ay = b + \Delta b$ with $\Delta b = -r$ exactly.
- The relative perturbation to the RHS is $\frac{\|r\|}{\|b\|}$.

Suppose we model entire perturbation in matrix A .

- Then y satisfies $(A + \frac{ry^T}{y^T y})y = b$
- The relative perturbation to the matrix is

$$\left\| \frac{ry^T}{y^T y} \right\|_2 / \|A\|_2 = \frac{\|r\|_2}{\|A\| \|y\|_2}$$

5-13

TB: 12; AB: 1.2.7; GvL 3.5 – PertA

5-13

Normwise backward error in both A & b

For a given y and given perturbation directions E, e_b , we define the Normwise backward error:

$$\eta_{E, e_b}(y) = \min\{\epsilon \mid (A + \Delta A)y = b + \Delta b; \\ \text{where } \Delta A, \Delta b \text{ satisfy: } \|\Delta A\| \leq \epsilon \|E\|; \\ \text{and } \|\Delta b\| \leq \epsilon \|e_b\|\}$$

In other words $\eta_{E, e_b}(y)$ is the smallest ϵ for which

$$(1) \begin{cases} (A + \Delta A)y = b + \Delta b; \\ \|\Delta A\| \leq \epsilon \|E\|; \quad \|\Delta b\| \leq \epsilon \|e_b\| \end{cases}$$

5-14

TB: 12; AB: 1.2.7; GvL 3.5 – PertA

5-14

- y is given (a computed solution). E and e_b to be selected (most likely 'directions of perturbation for A and b ').
- Typical choice: $E = A, e_b = b$

 Explain why this is not unreasonable

Let $r = b - Ay$. Then we have:

THEOREM 3: $\eta_{E, e_b}(y) = \frac{\|r\|}{\|E\| \|y\| + \|e_b\|}$


Normwise backward error is for case $E = A, e_b = b$:


$$\eta_{A, b}(y) = \frac{\|r\|}{\|A\| \|y\| + \|b\|}$$

5-15

TB: 12; AB: 1.2.7; GvL 3.5 – PertA

5-15

 Show how this can be used in practice as a means to stop some iterative method which computes a sequence of approximate solutions to $Ax = b$.

 Consider the 6×6 Vandermonde system $Ax = b$ where $a_{ij} = j^{2(i-1)}$, $b = A * [1, 1, \dots, 1]^T$. We perturb A by E , with $\|E\| \leq 10^{-10} \|A\|$ and b similarly and solve the system. Evaluate the backward error for this case. Evaluate the forward bound provided by Theorem 2. Comment on the results.

5-16

TB: 12; AB: 1.2.7; GvL 3.5 – PertA

5-16

Estimating condition numbers.

- Often we just want to get a lower bound for condition number [it is 'worse than ...']
- We want to estimate $\|A\| \|A^{-1}\|$.
- The norm $\|A\|$ is usually easy to compute but $\|A^{-1}\|$ is not.
- We want: Avoid the expense of computing A^{-1} explicitly.

Idea:

- Select a vector v so that $\|v\| = 1$ but $\|Av\| = \tau$ is small.
- Then: $\|A^{-1}\| \geq 1/\tau$ (show why) and:

$$\kappa(A) \geq \frac{\|A\|}{\tau}$$

5-17

TB: 12; AB: 1.2.8 ;GvL 3.5; Ort 9.3-4 – PertBshort

5-17

- Condition number worse than $\|A\|/\tau$.
- Typical choice for v : choose $[\dots \pm 1 \dots]$ with signs chosen on the fly during back-substitution to maximize the next entry in the solution, based on the upper triangular factor from Gaussian Elimination.
- Similar techniques used to estimate condition numbers of large matrices in matlab.

5-18

TB: 12; AB: 1.2.8 ;GvL 3.5; Ort 9.3-4 – PertBshort

5-18

Condition numbers and near-singularity

- $1/\kappa \approx$ relative distance to nearest singular matrix.

Let A, B be two $n \times n$ matrices with A nonsingular and B singular. Then

$$\frac{1}{\kappa(A)} \leq \frac{\|A - B\|}{\|A\|}$$

Proof: B singular $\rightarrow \exists x \neq 0$ such that $Bx = 0$.

$$\begin{aligned} \|x\| &= \|A^{-1}Ax\| \leq \|A^{-1}\| \|Ax\| = \|A^{-1}\| \|(A - B)x\| \\ &\leq \|A^{-1}\| \|A - B\| \|x\| \end{aligned}$$

Divide both sides by $\|x\| \times \kappa(A) = \|x\| \|A\| \|A^{-1}\|$ ➤ result. QED.

5-19

TB: 12; AB: 1.2.8 ;GvL 3.5; Ort 9.3-4 – PertBshort

5-19

Example:

$$\text{let } A = \begin{pmatrix} 1 & 1 \\ 1 & 0.99 \end{pmatrix} \text{ and } B = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

$$\text{Then } \frac{1}{\kappa_1(A)} \leq \frac{0.01}{2} \text{ ➤ } \kappa_1(A) \geq \frac{2}{0.01} = 200.$$

- It can be shown that (Kahan)

$$\frac{1}{\kappa(A)} = \min_B \left\{ \frac{\|A - B\|}{\|A\|} \mid \det(B) = 0 \right\}$$

5-20

TB: 12; AB: 1.2.8 ;GvL 3.5; Ort 9.3-4 – PertBshort

5-20

Estimating errors from residual norms

Let \tilde{x} an approximate solution to system $Ax = b$ (e.g., computed from an iterative process). We can compute the residual norm:

$$\|r\| = \|b - A\tilde{x}\|$$

Question: How to estimate the error $\|x - \tilde{x}\|$ from $\|r\|$?

- One option is to use the inequality

$$\frac{\|x - \tilde{x}\|}{\|x\|} \leq \kappa(A) \frac{\|r\|}{\|b\|}.$$

- We must have an estimate of $\kappa(A)$.

Proof of inequality.

First, note that $A(x - \tilde{x}) = b - A\tilde{x} = r$. So:

$$\|x - \tilde{x}\| = \|A^{-1}r\| \leq \|A^{-1}\| \|r\|$$

Also note that from the relation $b = Ax$, we get

$$\|b\| = \|Ax\| \leq \|A\| \|x\| \rightarrow \|x\| \geq \frac{\|b\|}{\|A\|}$$

Therefore,

$$\frac{\|x - \tilde{x}\|}{\|x\|} \leq \frac{\|A^{-1}\| \|r\|}{\|b\|/\|A\|} = \kappa(A) \frac{\|r\|}{\|b\|} \quad \square$$

 Show that

$$\frac{\|x - \tilde{x}\|}{\|x\|} \geq \frac{1}{\kappa(A)} \frac{\|r\|}{\|b\|}.$$