

Least-Squares Systems and The QR factorization

- Orthogonality
- Least-squares systems.
- The Gram-Schmidt and Modified Gram-Schmidt processes.
- The Householder QR and the Givens QR.

7-1

Orthogonality

1. Two vectors u and v are orthogonal if $(u, v) = 0$.
 2. A system of vectors $\{v_1, \dots, v_n\}$ is **orthogonal** if $(v_i, v_j) = 0$ for $i \neq j$; and **orthonormal** if $(v_i, v_j) = \delta_{ij}$
 3. A matrix is **orthogonal** if its columns are orthonormal
- Notation: $V = [v_1, \dots, v_n] \implies$ matrix with column-vectors v_1, \dots, v_n .
 - Orthogonality is essential in understanding and solving least-squares problems.

7-2

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-2

Least-Squares systems

- Given: an $m \times n$ matrix $n < m$. Problem: find x which minimizes:

$$\|b - Ax\|_2$$

- Good illustration: Data fitting.

Typical problem of data fitting: We seek an unknown function as a linear combination ϕ of n known functions ϕ_i (e.g. polynomials, trig. functions). Experimental data (not accurate) provides measures β_1, \dots, β_m of this unknown function at points t_1, \dots, t_m . Problem: find the 'best' possible approximation ϕ to this data.

$$\phi(t) = \sum_{i=1}^n \xi_i \phi_i(t) \quad , \quad \text{s.t.} \quad \phi(t_j) \approx \beta_j, \quad j = 1, \dots, m$$

7-3

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-3

- Question: Close in what sense?
- Least-squares approximation: Find ϕ such that

$$\phi(t) = \sum_{i=1}^n \xi_i \phi_i(t), \quad \& \quad \sum_{j=1}^m |\phi(t_j) - \beta_j|^2 = \text{Min}$$

- In linear algebra terms: find 'best' approximation to a vector b from linear combinations of vectors $f_i, i = 1, \dots, n$, where

$$b = \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_m \end{pmatrix}, \quad f_i = \begin{pmatrix} \phi_i(t_1) \\ \phi_i(t_2) \\ \vdots \\ \phi_i(t_m) \end{pmatrix}$$

7-4

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-4

► We want to find $x = \{\xi_i\}_{i=1,\dots,n}$ such that


$$\left\| \sum_{i=1}^n \xi_i f_i - b \right\|_2 \quad \text{Minimum}$$

Define

$$F = [f_1, f_2, \dots, f_n], \quad x = \begin{pmatrix} \xi_1 \\ \vdots \\ \xi_n \end{pmatrix}$$

► We want to find x to **minimize $\|b - Fx\|_2$**

► This is a **Least-squares linear system**: F is $m \times n$, with $m \geq n$.

 Formulate the least-squares system for the problem of finding the polynomial of degree 2 that approximates a function f which satisfies $f(-1) = -1; f(0) = 1; f(1) = 2; f(2) = 0$

Solution: $\phi_1(t) = 1; \phi_2(t) = t; \phi_3(t) = t^2;$

• Evaluate the ϕ_i 's at points $t_1 = -1; t_2 = 0; t_3 = 1; t_4 = 2$:

$$f_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \quad f_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \\ 2 \end{pmatrix} \quad f_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 4 \end{pmatrix} \quad \rightarrow$$

► So the coefficients ξ_1, ξ_2, ξ_3 of the polynomial $\xi_1 + \xi_2 t + \xi_3 t^2$ are the solution of the least-squares problem $\min \|b - Fx\|$ where:

$$F = \begin{pmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{pmatrix} \quad b = \begin{pmatrix} -1 \\ 1 \\ 2 \\ 0 \end{pmatrix}$$

THEOREM. The vector x_* minimizes $\psi(x) = \|b - Fx\|_2^2$ if and only if it is the solution of the **normal equations**:

$$F^T F x = F^T b$$

Proof: Expand out the formula for $\psi(x_* + \delta x)$:

$$\begin{aligned} \psi(x_* + \delta x) &= ((b - Fx_*) - F\delta x)^T ((b - Fx_*) - F\delta x) \\ &= \psi(x_*) - 2(F\delta x)^T (b - Fx_*) + (F\delta x)^T (F\delta x) \\ &= \psi(x_*) - 2(\delta x)^T \underbrace{[F^T (b - Fx_*)]}_{-\nabla_x \psi} + \underbrace{(F\delta x)^T (F\delta x)}_{\text{always positive}} \end{aligned}$$

Can see that $\psi(x_* + \delta x) \geq \psi(x_*)$ for any δx , iff the boxed quantity [the gradient vector] is zero. Q.E.D.





Illustration of theorem: x^* is the best approximation to the vector b from the subspace $\text{span}\{F\}$ if and only if $b - Fx^*$ is \perp to the whole subspace $\text{span}\{F\}$. This in turn is equivalent to $F^T (b - Fx^*) = 0$ ► Normal equations.

Example:

Points:	$t_1 = -1$	$t_2 = -1/2$	$t_3 = 0$	$t_4 = 1/2$	$t_5 = 1$
Values:	$\beta_1 = 0.1$	$\beta_2 = 0.3$	$\beta_3 = 0.3$	$\beta_4 = 0.2$	$\beta_5 = 0.0$



7-9 TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-9

1) Approximations by polynomials of degree one:

➤ $\phi_1(t) = 1, \phi_2(t) = t$.


$$F = \begin{pmatrix} 1.0 & -1.0 \\ 1.0 & -0.5 \\ 1.0 & 0 \\ 1.0 & 0.5 \\ 1.0 & 1.0 \end{pmatrix}$$

$$F^T F = \begin{pmatrix} 5.0 & 0 \\ 0 & 2.5 \end{pmatrix}$$

$$F^T b = \begin{pmatrix} 0.9 \\ -0.15 \end{pmatrix}$$

➤ Best approximation is

$$\phi(t) = 0.18 - 0.06t$$



7-10

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR


7-10

2) Approximation by polynomials of degree 2:

➤ $\phi_1(t) = 1, \phi_2(t) = t, \phi_3(t) = t^2$.

➤ Best polynomial found:

$$0.3085714285 - 0.06 \times t - 0.2571428571 \times t^2$$



7-11 TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-11

Problem with Normal Equations

➤ Condition number is high: if A is square and non-singular, then

$$\kappa_2(A) = \|A\|_2 \cdot \|A^{-1}\|_2 = \sigma_{\max}/\sigma_{\min}$$

$$\kappa_2(A^T A) = \|A^T A\|_2 \cdot \|(A^T A)^{-1}\|_2 = (\sigma_{\max}/\sigma_{\min})^2$$

➤ Example: Let $A = \begin{pmatrix} 1 & 1 & -\epsilon \\ \epsilon & 0 & 1 \\ 0 & \epsilon & 1 \end{pmatrix}$.

➤ Then $\kappa(A) \approx \sqrt{2}/\epsilon$, but $\kappa(A^T A) \approx 2\epsilon^{-2}$.

$$\text{fl}(A^T A) = \text{fl} \begin{pmatrix} 2 + \epsilon^2 & 1 & 0 \\ 1 & 1 + \epsilon^2 & 0 \\ 0 & 0 & 1 + \epsilon^2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

is singular to working precision (if $\epsilon < \underline{u}$).

7-12

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-12

Finding an orthonormal basis of a subspace

- Goal: Find vector in $\text{span}(X)$ closest to b .
- Much easier with an orthonormal basis for $\text{span}(X)$.

Problem: Given $X = [x_1, \dots, x_n]$, compute $Q = [q_1, \dots, q_n]$ which has orthonormal columns and s.t. $\text{span}(Q) = \text{span}(X)$

- Note: each column of X must be a linear combination of certain columns of Q .
- We will find Q so that x_j (j column of X) is a linear combination of the first j columns of Q .


7-13

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-13

ALGORITHM : 1. Classical Gram-Schmidt

1. For $j = 1, \dots, n$ Do:
2. Set $\hat{q} := x_j$
3. Compute $r_{ij} := (\hat{q}, q_i)$, for $i = 1, \dots, j - 1$
4. For $i = 1, \dots, j - 1$ Do :
5. Compute $\hat{q} := \hat{q} - r_{ij}q_i$
6. EndDo
7. Compute $r_{jj} := \|\hat{q}\|_2$,
8. If $r_{jj} = 0$ then Stop, else $q_j := \hat{q}/r_{jj}$
9. EndDo

- All n steps can be completed iff x_1, x_2, \dots, x_n are linearly independent.  Prove this result

7-14

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-14

- Lines 5 and 7-8 show that

$$x_j = r_{1j}q_1 + r_{2j}q_2 + \dots + r_{jj}q_j$$

- If $X = [x_1, x_2, \dots, x_n]$, $Q = [q_1, q_2, \dots, q_n]$, and if R is the $n \times n$ upper triangular matrix

$$R = \{r_{ij}\}_{i,j=1,\dots,n}$$

then the above relation can be written as

$$X = QR$$

- R is upper triangular, Q is orthogonal. This is called the QR factorization of X .

 What is the cost of the factorization when $X \in \mathbb{R}^{m \times n}$?

7-15

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-15

$$\begin{array}{c}
 \boxed{X} \\
 \text{Original} \\
 \text{matrix}
 \end{array}
 =
 \begin{array}{c}
 \boxed{Q} \\
 \text{Q is orthogonal} \\
 (Q^H Q = I)
 \end{array}
 *
 \begin{array}{c}
 \boxed{R} \\
 \text{R is upper} \\
 \text{triangular}
 \end{array}$$

Another decomposition:

A matrix X , with linearly independent columns, is the product of an orthogonal matrix Q and a upper triangular matrix R .

7-16

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-16

- Better algorithm: Modified Gram-Schmidt.

ALGORITHM : 2. Modified Gram-Schmidt

1. For $j = 1, \dots, n$ Do:
2. Define $\hat{q} := x_j$
3. For $i = 1, \dots, j - 1$, Do:
4. $r_{ij} := (\hat{q}, q_i)$
5. $\hat{q} := \hat{q} - r_{ij}q_i$
6. EndDo
7. Compute $r_{jj} := \|\hat{q}\|_2$,
8. If $r_{jj} = 0$ then Stop, else $q_j := \hat{q}/r_{jj}$
9. EndDo

Only difference: inner product uses the accumulated subsum instead of original \hat{q}

7-17


TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-17

The operations in lines 4 and 5 can be written as

$$\hat{q} := ORTH(\hat{q}, q_i)$$

where $ORTH(x, q)$ denotes the operation of orthogonalizing a vector x against a unit vector q .



Result of $z = ORTH(x, q)$

7-18

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-18

- Modified Gram-Schmidt algorithm is much more stable than classical Gram-Schmidt in general. [A few examples easily show this].

Suppose MGS is applied to A yielding computed matrices \hat{Q} and \hat{R} . Then there are constants c_i (depending on (m, n)) such that

$$A + E_1 = \hat{Q}\hat{R} \quad \|E_1\|_2 \leq c_1 \underline{u} \|A\|_2$$

$$\|\hat{Q}^T \hat{Q} - I\|_2 \leq c_2 \underline{u} \kappa_2(A) + O((\underline{u} \kappa_2(A))^2)$$

for a certain perturbation matrix E_1 , and there exists an orthonormal matrix Q such that

$$A + E_2 = Q\hat{R} \quad \|E_2(:, j)\|_2 \leq c_3 \underline{u} \|A(:, j)\|_2$$

for a certain perturbation matrix E_2 .

7-19

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-19

- An equivalent version:

ALGORITHM : 3. Modified Gram-Schmidt - 2 -

0. Set $\hat{Q} := X$
1. For $i = 1, \dots, n$ Do:
2. Compute $r_{ii} := \|\hat{q}_i\|_2$,
3. If $r_{ii} = 0$ then Stop, else $q_i := \hat{q}_i/r_{ii}$
4. For $j = i + 1, \dots, n$, Do:
5. $r_{ij} := (\hat{q}_j, q_i)$
6. $\hat{q}_j := \hat{q}_j - r_{ij}q_i$
7. EndDo
8. EndDo

- Does exactly the same computation as previous algorithm, but in a different order.

7-20

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-20

Example:

Orthonormalize the system of vectors:

$$X = [x_1, x_2, x_3] = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & -1 \\ 1 & 0 & 4 \end{pmatrix}$$

Answer:

$$q_1 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}; \quad \hat{q}_2 = x_2 - (x_2, q_1)q_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} - 1 \times \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}$$

$$\hat{q}_2 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}; \quad q_2 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}$$

$$\hat{q}_3 = x_3 - (x_3, q_1)q_1 = \begin{pmatrix} 1 \\ 0 \\ -1 \\ 4 \end{pmatrix} - 2 \times \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \\ -2 \\ 3 \end{pmatrix}$$

$$\hat{q}_3 = \hat{q}_3 - (\hat{q}_3, q_2)q_2 = \begin{pmatrix} 0 \\ -1 \\ -2 \\ 3 \end{pmatrix} - (-1) \times \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \\ -2.5 \\ 2.5 \end{pmatrix}$$

$$\|\hat{q}_3\|_2 = \sqrt{13} \rightarrow q_3 = \frac{\hat{q}_3}{\|\hat{q}_3\|_2} = \frac{1}{\sqrt{13}} \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \\ -2.5 \\ 2.5 \end{pmatrix}$$

 For this example: what is Q ? what is R ? Compute $Q^T Q$.

➤ Result is the identity matrix.

Recall: For any orthogonal matrix Q , we have

$$Q^T Q = I$$

(In complex case: $Q^H Q = I$).

Consequence: For an $n \times n$ orthogonal matrix $Q^{-1} = Q^T$.
(Q is orthogonal/ unitary)

Application: another method for solving linear systems.

$$Ax = b$$

A is an $n \times n$ nonsingular matrix. Compute its QR factorization.

➤ Multiply both sides by $Q^T \rightarrow Q^T Q R x = Q^T b \rightarrow$

$$R x = Q^T b$$

Method:

➤ Compute the QR factorization of A , $A = QR$.

➤ Solve the upper triangular system $R x = Q^T b$.

 Cost??

Use of the QR factorization

Problem: $Ax \approx b$ in least-squares sense

A is an $m \times n$ (full-rank) matrix. Let

$$A = QR$$

the QR factorization of A and consider the normal equations:

$$A^T Ax = A^T b \rightarrow R^T Q^T QRx = R^T Q^T b \rightarrow$$

$$R^T Rx = R^T Q^T b \rightarrow Rx = Q^T b$$

(R^T is an $n \times n$ nonsingular matrix). Therefore,

$$x = R^{-1} Q^T b$$

7-25

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-25

Another derivation:

- Recall: $\text{span}(Q) = \text{span}(A)$
- So $\|b - Ax\|_2$ is minimum when $b - Ax \perp \text{span}\{Q\}$
- Therefore solution x must satisfy $Q^T(b - Ax) = 0 \rightarrow$

$$Q^T(b - QRx) = 0 \rightarrow Rx = Q^T b$$

$$x = R^{-1} Q^T b$$

7-26

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-26

- Also observe that for any vector w

$$w = QQ^T w + (I - QQ^T)w$$

and that $w = QQ^T w \perp (I - QQ^T)w \rightarrow$

- Pythagoras theorem $\rightarrow \|w\|_2^2 = \|QQ^T w\|_2^2 + \|(I - QQ^T)w\|_2^2$

$$\begin{aligned} \|b - Ax\|^2 &= \|b - QRx\|^2 \\ &= \|(I - QQ^T)b + Q(Q^T b - Rx)\|^2 \\ &= \|(I - QQ^T)b\|^2 + \|Q(Q^T b - Rx)\|^2 \\ &= \|(I - QQ^T)b\|^2 + \|Q^T b - Rx\|^2 \end{aligned}$$

- Min is reached when 2nd term of r.h.s. is zero.

7-27


TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR


7-27

Method:

- Compute the QR factorization of A , $A = QR$.
- Compute the right-hand side $f = Q^T b$
- Solve the upper triangular system $Rx = f$.
- x is the least-squares solution

- As a rule it is not a good idea to form $A^T A$ and solve the normal equations. Methods using the QR factorization are better.

 6 Total cost?? (depends on the algorithm used to get the QR decomposition).

 7 Using matlab find the parabola that fits the data in previous data fitting example (p. 8-10) in L.S. sense [verify that the result found is correct.]

7-28

TB: 7,8,11,19; AB: 2.1, 2.3.4, ;GvL 5, 5.3 – QR

7-28