



Class time : MW 4:00 – 5:15 pm
Room : Keller 3-230 or Online
Instructor : Daniel Boley

Lecture notes: <http://www-users.cselabs.umn.edu/classes/Fall-2021/csci5304/>

August 27, 2021

- The Pseudo-inverse
- Use of SVD for least-squares problems
- Application to regularization
- Numerical rank

10-1

Pseudo-inverse of an arbitrary matrix

➤ Let $A = U\Sigma V^T$ which we rewrite as

$$A = (U_1 \ U_2) \begin{pmatrix} \Sigma_1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} V_1^T \\ V_2^T \end{pmatrix} = U_1 \Sigma_1 V_1^T$$

Then the pseudo inverse of A is

$$A^\dagger = V_1 \Sigma_1^{-1} U_1^T = \sum_{j=1}^r \frac{1}{\sigma_j} v_j u_j^T$$

➤ The pseudo-inverse of A is the mapping from a vector b to the solution $\min_x \|Ax - b\|_2^2$ that has minimal norm (to be shown)

➤ In the full-rank overdetermined case, the normal equations yield $x = \underbrace{(A^T A)^{-1} A^T}_{A^\dagger} b$

Least-squares problem via the SVD

Pb: $\min \|b - Ax\|_2$ in general case. Consider SVD of A :

$$A = (U_1 \ U_2) \begin{pmatrix} \Sigma_1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} V_1^T \\ V_2^T \end{pmatrix} = \sum_{i=1}^r \sigma_i v_i u_i^T$$


➤ Find all possible least-squares solutions. Also find the one with min. 2-norm.

1) Express x in V basis : $x = Vy = [V_1, V_2] \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}$

2) Then left multiply by U^T to get

$$\|Ax - b\|_2^2 = \left\| \begin{pmatrix} \Sigma_1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} - \begin{pmatrix} U_1^T b \\ U_2^T b \end{pmatrix} \right\|_2^2$$

3) Find all possible solutions in terms of $y = [y_1; y_2]$

 What are **all** least-squares solutions to the above system? Among these which one has minimum norm?

10-3

GvL 2.4, 5.4-5 – SVD1

10-3

Answer: From above, must have $y_1 = \Sigma_1^{-1} U_1^T b$ and $y_2 =$ anything (free).

Recall that:

$$\begin{aligned} x &= [V_1, V_2] \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = V_1 y_1 + V_2 y_2 \\ &= V_1 \Sigma_1^{-1} U_1^T b + V_2 y_2 \\ &= \boxed{A^\dagger b + V_2 y_2} \end{aligned}$$

➤ Note: $A^\dagger b \in \text{Ran}(A^T)$ and $V_2 y_2 \in \text{Null}(A)$.

➤ Therefore: least-squares solutions are all of the form $A^\dagger b + w$ where $w \in \text{Null}(A)$.

➤ Smallest norm when $y_2 = 0$.


10-4

GvL 2.4, 5.4-5 – SVD1

10-4

➤ Minimum norm solution to $\min_x \|Ax - b\|_2^2$ satisfies $\Sigma_1 y_1 = U_1^T b$, $y_2 = 0$. It is:

$$x_{LS} = V_1 \Sigma_1^{-1} U_1^T b = A^\dagger b$$

 If $A \in \mathbb{R}^{m \times n}$ what are the dimensions of A^\dagger , $A^\dagger A$, AA^\dagger ?

 Show that $A^\dagger A$ is an orthogonal projector. What are its range and null-space?

 Same questions for AA^\dagger .

10-5

GvL 2.4, 5.4-5 – SVD1

10-5

Moore-Penrose Inverse

The pseudo-inverse of A is given by

$$A^\dagger = V \begin{pmatrix} \Sigma_1^{-1} & 0 \\ 0 & 0 \end{pmatrix} U^T = \sum_{i=1}^r \frac{v_i u_i^T}{\sigma_i}$$

Moore-Penrose conditions:

The pseudo inverse of a matrix is uniquely determined by these four conditions:

- (1) $AXA = A$
- (2) $XAX = X$
- (3) $(AX)^H = AX$
- (4) $(XA)^H = XA$

➤ In the full-rank overdetermined case, $A^\dagger = (A^T A)^{-1} A^T$

10-6

GvL 2.4, 5.4-5 – SVD1

10-6

Least-squares problems and the SVD

- The SVD can give much information on solutions of overdetermined and underdetermined linear systems.

Let A be an $m \times n$ matrix and $A = U\Sigma V^T$ its SVD with $r = \text{rank}(A)$, $V = [v_1, \dots, v_n]$ $U = [u_1, \dots, u_m]$. Then

$$x_{LS} = \sum_{i=1}^r \frac{u_i^T b}{\sigma_i} v_i$$

minimizes $\|b - Ax\|_2$ and has the smallest 2-norm among all possible minimizers. In addition,

$$\rho_{LS} \equiv \|b - Ax_{LS}\|_2 = \|z\|_2 \text{ with } z = [u_{r+1}, \dots, u_m]^T b$$

10-7

GvL 2.4, 5.4-5 – SVD1

10-7

Least-squares problems and pseudo-inverses

- A restatement of the first part of the previous result:

Consider the general linear least-squares problem

$$\min_{x \in S} \|x\|_2, \quad S = \{x \in \mathbb{R}^n \mid \|b - Ax\|_2 \text{ min}\}.$$

This problem always has a unique solution given by

$$x = A^\dagger b$$

10-8

GvL 2.4, 5.4-5 – SVD1

10-8

10-5 Consider the matrix:

$$A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 0 & -2 & 1 \end{pmatrix}$$

- Compute the thin SVD of A
- Find the matrix B of rank 1 which is the closest to the above matrix in the 2-norm sense.
- What is the pseudo-inverse of A ?
- What is the pseudo-inverse of B ?
- Find the vector x of smallest norm which minimizes $\|b - Ax\|_2$ with $b = (1, 1)^T$
- Find the vector x of smallest norm which minimizes $\|b - Bx\|_2$ with $b = (1, 1)^T$

10-9

GvL 2.4, 5.4-5 – SVD1

10-9

Ill-conditioned systems and the SVD

- Let A be $m \times m$ and $A = U\Sigma V^T$ its SVD
- Solution of $Ax = b$ is $x = A^{-1}b = \sum_{i=1}^m \frac{u_i^T b}{\sigma_i} v_i$
- When A is very ill-conditioned, it has many small singular values. The division by these small σ_i 's will amplify any noise in the data. If $\tilde{b} = b + \epsilon$ then

$$A^{-1}\tilde{b} = \sum_{i=1}^m \frac{u_i^T b}{\sigma_i} v_i + \underbrace{\sum_{i=1}^m \frac{u_i^T \epsilon}{\sigma_i} v_i}_{\text{Error}}$$

- Result: solution could be completely meaningless.

10-10

GvL 2.4, 5.4-5 – SVD1

10-10

Remedy: SVD regularization

Truncate the SVD by only keeping the σ_i 's that are $\geq \tau$, where τ is a threshold

- Gives the Truncated SVD solution (TSVD solution:)

$$x_{TSVD} = \sum_{\sigma_i \geq \tau} \frac{u_i^T b}{\sigma_i} v_i$$

- Many applications [e.g., Image and signal processing,...]

Numerical rank and the SVD

- Assuming the original matrix A is exactly of rank k the **computed** SVD of A will be the SVD of a nearby matrix $A + E$ – Can show: $|\hat{\sigma}_i - \sigma_i| \leq \alpha \sigma_{i+1}$

- Result: zero singular values will yield small computed singular values and r larger sing. values.

- Reverse problem: *numerical rank* – The ϵ -rank of A :

$$r_\epsilon = \min\{\text{rank}(B) : B \in \mathbb{R}^{m \times n}, \|A - B\|_2 \leq \epsilon\},$$

- ☒ Show that r_ϵ equals the number sing. values that are $> \epsilon$

- ☒ Show: r_ϵ equals the number of columns of A that are linearly independent for any perturbation of A with norm $\leq \epsilon$.

- Practical problem : How to set ϵ ?

Pseudo inverses of full-rank matrices

Case 1: $m \geq n$ Then $A^\dagger = (A^T A)^{-1} A^T$

- Thin SVD is $A = U_1 \Sigma_1 V_1^T$ and V_1, Σ_1 are $n \times n$. Then:

$$\begin{aligned} (A^T A)^{-1} A^T &= (V_1 \Sigma_1^2 V_1^T)^{-1} V_1 \Sigma_1 U_1^T \\ &= V_1 \Sigma_1^{-2} V_1^T V_1 \Sigma_1 U_1^T \\ &= V_1 \Sigma_1^{-1} U_1^T \\ &= A^\dagger \end{aligned}$$

Example: Pseudo-inverse of $\begin{pmatrix} 0 & 1 \\ 1 & 2 \\ 2 & -1 \\ 0 & 1 \end{pmatrix}$ is?

Case 2: $m < n$ Then $A^\dagger = A^T (A A^T)^{-1}$

- Thin SVD is $A = U_1 \Sigma_1 V_1^T$. Now U_1, Σ_1 are $m \times m$ and:

$$\begin{aligned} A^T (A A^T)^{-1} &= V_1 \Sigma_1 U_1^T [U_1 \Sigma_1^2 U_1^T]^{-1} \\ &= V_1 \Sigma_1 U_1^T U_1 \Sigma_1^{-2} U_1^T \\ &= V_1 \Sigma_1^{-1} U_1^T \\ &= A^\dagger \end{aligned}$$

Example: Pseudo-inverse of $\begin{pmatrix} 0 & 1 & 2 & 0 \\ 1 & 2 & -1 & 1 \end{pmatrix}$ is?

- Mnemonic: The pseudo inverse of A is A^T completed by the inverse of the smaller of $(A^T A)^{-1}$ or $(A A^T)^{-1}$ where it fits (i.e., left or right)