

THE URV & SINGULAR VALUE DECOMPOSITIONS

- Orthogonal subspaces;
- Orthogonal projectors; Orthogonal decomposition;
- The URV decomposition
- Introduction to the Singular Value Decomposition
- The SVD – existence and properties.

9-1

Orthogonal projectors and subspaces

Notation: Given a subspace \mathcal{X} of \mathbb{R}^m define

$$\mathcal{X}^\perp = \{y \mid y \perp x, \forall x \in \mathcal{X}\}$$

➤ Let $Q = [q_1, \dots, q_r]$ an orthonormal basis of \mathcal{X}

🔗 How would you obtain such a basis?

➤ Then define orthogonal projector $P = QQ^T$

Properties

- (a) $P^2 = P$ (b) $(I - P)^2 = I - P$
 (c) $\text{Ran}(P) = \mathcal{X}$ (d) $\text{Null}(P) = \mathcal{X}^\perp$
 (e) $\text{Ran}(I - P) = \text{Null}(P) = \mathcal{X}^\perp$

➤ Note that (b) means that $I - P$ is also a projector

9-2

AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-2

Proof. (a), (b) are trivial

(c): Clearly $\text{Ran}(P) = \{x \mid x = QQ^T y, y \in \mathbb{R}^m\} \subseteq \mathcal{X}$.
 Any $x \in \mathcal{X}$ is of the form $x = Qy, y \in \mathbb{R}^m$. Take $Px = QQ^T(Qy) = Qy = x$. Since $x = Px, x \in \text{Ran}(P)$. So $\mathcal{X} \subseteq \text{Ran}(P)$. In the end $\mathcal{X} = \text{Ran}(P)$.

(d): $x \in \mathcal{X}^\perp \Leftrightarrow (x, y) = 0, \forall y \in \mathcal{X} \Leftrightarrow (x, Qz) = 0, \forall z \in \mathbb{R}^r \Leftrightarrow (Q^T x, z) = 0, \forall z \in \mathbb{R}^r \Leftrightarrow Q^T x = 0 \Leftrightarrow QQ^T x = 0 \Leftrightarrow Px = 0$.

(e): Need to show inclusion both ways.

• $x \in \text{Null}(P) \Leftrightarrow Px = 0 \Leftrightarrow (I - P)x = x \rightarrow x \in \text{Ran}(I - P)$

• $x \in \text{Ran}(I - P) \Leftrightarrow \exists y \in \mathbb{R}^m \mid x = (I - P)y \rightarrow Px = P(I - P)y = 0 \rightarrow x \in \text{Null}(P)$ \square

9-3

AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-3

Result: Any $x \in \mathbb{R}^m$ can be written in a unique way as

$$x = x_1 + x_2, \quad x_1 \in \mathcal{X}, \quad x_2 \in \mathcal{X}^\perp$$

➤ Proof: Just set $x_1 = Px, \quad x_2 = (I - P)x$

➤ Note:

$$\mathcal{X} \cap \mathcal{X}^\perp = \{0\}$$

➤ Therefore:

$$\mathbb{R}^m = \mathcal{X} \oplus \mathcal{X}^\perp$$

➤ Called the **Orthogonal Decomposition**

9-4

AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-4

Orthogonal decomposition

- In other words $\mathbb{R}^m = P\mathbb{R}^m \oplus (I - P)\mathbb{R}^m$ or:
 $\mathbb{R}^m = \text{Ran}(P) \oplus \text{Ran}(I - P)$ or:
 $\mathbb{R}^m = \text{Ran}(P) \oplus \text{Null}(P)$ or:
 $\mathbb{R}^m = \text{Ran}(P) \oplus \text{Ran}(P)^\perp$
- Can complete basis $\{q_1, \dots, q_r\}$ into orthonormal basis of \mathbb{R}^m , q_{r+1}, \dots, q_m
- $\{q_{r+1}, \dots, q_m\} = \text{basis of } \mathcal{X}^\perp. \rightarrow \text{dim}(\mathcal{X}^\perp) = m - r.$

Four fundamental subspaces - URV decomposition

Let $A \in \mathbb{R}^{m \times n}$ and consider $\text{Ran}(A)^\perp$

Property 1: $\text{Ran}(A)^\perp = \text{Null}(A^T)$

Proof: $x \in \text{Ran}(A)^\perp$ iff $(Ay, x) = 0$ for all y iff $(y, A^T x) = 0$ for all y ...

Property 2: $\text{Ran}(A^T) = \text{Null}(A)^\perp$

- Take $\mathcal{X} = \text{Ran}(A)$ in orthogonal decomposition. ➤ Result:

$$\begin{aligned} \mathbb{R}^m &= \text{Ran}(A) \oplus \text{Null}(A^T) \\ \mathbb{R}^n &= \text{Ran}(A^T) \oplus \text{Null}(A) \end{aligned}$$

4 fundamental subspaces
 $\text{Ran}(A)$ $\text{Null}(A^T)$
 $\text{Ran}(A^T)$ $\text{Null}(A)$

- Express the above with bases for \mathbb{R}^m :

$$\underbrace{[u_1, u_2, \dots, u_r]}_{\text{Ran}(A)} \underbrace{[u_{r+1}, u_{r+2}, \dots, u_m]}_{\text{Null}(A^T)}$$

and for \mathbb{R}^n $\underbrace{[v_1, v_2, \dots, v_r]}_{\text{Ran}(A^T)} \underbrace{[v_{r+1}, v_{r+2}, \dots, v_n]}_{\text{Null}(A)}$

- Observe $u_i^T A v_j = 0$ for $i > r$ or $j > r$. Therefore

$$U^T A V = R = \begin{pmatrix} C & 0 \\ 0 & 0 \end{pmatrix}_{m \times n} \quad C \in \mathbb{R}^{r \times r} \rightarrow$$


$$A = URV^T$$

- General class of URV decompositions

- Far from unique.

 Show how you can get a decomposition in which C is lower (or upper) triangular, from the above factorization.

- Can select decomposition so that R is upper triangular \rightarrow URV decomposition.
- Can select decomposition so that R is lower triangular \rightarrow ULV decomposition.
- SVD = special case of URV where R = diagonal

 How can you get the ULV decomposition by using only the Householder QR factorization (possibly with pivoting)? [Hint: you must use Householder twice]

The Singular Value Decomposition (SVD)

Theorem For any matrix $A \in \mathbb{R}^{m \times n}$ there exist unitary matrices $U \in \mathbb{R}^{m \times m}$ and $V \in \mathbb{R}^{n \times n}$ such that

$$A = U \Sigma V^T$$

where Σ is a diagonal matrix with entries $\sigma_{ii} \geq 0$.

$$\sigma_{11} \geq \sigma_{22} \geq \dots \geq \sigma_{pp} \geq 0 \text{ with } p = \min(n, m)$$

► The σ_{ii} 's are the **singular values**. Notation change $\sigma_{ii} \rightarrow \sigma_i$

Proof: Let $\sigma_1 = \|A\|_2 = \max_{x, \|x\|_2=1} \|Ax\|_2$. There exists a pair of unit vectors v_1, u_1 such that

$$Av_1 = \sigma_1 u_1$$

9-9 AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-9

► Complete v_1 into an orthonormal basis of \mathbb{R}^n

$$V \equiv [v_1, V_2] = n \times n \text{ unitary}$$

► Complete u_1 into an orthonormal basis of \mathbb{R}^m

$$U \equiv [u_1, U_2] = m \times m \text{ unitary}$$

 Define U, V as single Householder reflectors.

► Then, it is easy to show that

$$AV = U \times \begin{pmatrix} \sigma_1 & w^T \\ 0 & B \end{pmatrix} \rightarrow U^T AV = \begin{pmatrix} \sigma_1 & w^T \\ 0 & B \end{pmatrix} \equiv A_1$$

9-10 AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-10

► Observe that

$$\left\| A_1 \begin{pmatrix} \sigma_1 \\ w \end{pmatrix} \right\|_2 \geq \sigma_1^2 + \|w\|^2 = \sqrt{\sigma_1^2 + \|w\|^2} \left\| \begin{pmatrix} \sigma_1 \\ w \end{pmatrix} \right\|_2$$


► This shows that w must be zero [why?]

► Complete the proof by an induction argument. ■


9-11 AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-11

Case 1:



Case 2:



9-12 AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-12

The “thin” SVD

- Consider the Case-1. It can be rewritten as

$$A = [U_1 U_2] \begin{pmatrix} \Sigma_1 \\ 0 \end{pmatrix} V^T$$

Which gives:

$$A = U_1 \Sigma_1 V^T$$

where U_1 is $m \times n$ (same shape as A), and Σ_1 and V are $n \times n$

- Referred to as the “thin” SVD. Important in practice.

 How can you obtain the thin SVD from the QR factorization of A and the SVD of an $n \times n$ matrix?

9-13 AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-13

A few properties. Assume that

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r > 0 \text{ and } \sigma_{r+1} = \dots = \sigma_p = 0$$

Then:

- $\text{rank}(A) = r =$ number of nonzero singular values.
- $\text{Ran}(A) = \text{span}\{u_1, u_2, \dots, u_r\}$
- $\text{Null}(A^T) = \text{span}\{u_{r+1}, u_{r+2}, \dots, u_m\}$
- $\text{Ran}(A^T) = \text{span}\{v_1, v_2, \dots, v_r\}$
- $\text{Null}(A) = \text{span}\{v_{r+1}, v_{r+2}, \dots, v_n\}$

9-14 AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-14

Properties of the SVD (continued)

- The matrix A admits the SVD expansion:

$$A = \sum_{i=1}^r \sigma_i u_i v_i^T$$

- $\|A\|_2 = \sigma_1 =$ largest singular value
- $\|A\|_F = (\sum_{i=1}^r \sigma_i^2)^{1/2}$
- When A is an $n \times n$ nonsingular matrix then $\|A^{-1}\|_2 = 1/\sigma_n$

Theorem Let $k < r$ and

$$A_k = \sum_{i=1}^k \sigma_i u_i v_i^T$$

then

$$\min_{\text{rank}(B)=k} \|A - B\|_2 = \|A - A_k\|_2 = \sigma_{k+1}$$

9-16 AB: 1.1. 2.2. 2.4; TB: 4-5; GvL 2.4, 5.4-5 – SVD

9-15

9-16

Proof: First: $\|A - B\|_2 \geq \sigma_{k+1}$, for any rank- k matrix B .

Consider $\mathcal{X} = \text{span}\{v_1, v_2, \dots, v_{k+1}\}$. Note:

$$\dim(\text{Null}(B)) = n - k \rightarrow \text{Null}(B) \cap \mathcal{X} \neq \{0\}$$

[Why?]

Let $x_0 \in \text{Null}(B) \cap \mathcal{X}$, $x_0 \neq 0$. Write $x_0 = Vy$. Then

$$\|(A - B)x_0\|_2 = \|Ax_0\|_2 = \|U\Sigma V^T Vy\|_2 = \|\Sigma y\|_2$$

But $\|\Sigma y\|_2 \geq \sigma_{k+1}\|x_0\|_2$ (Show this). $\rightarrow \|A - B\|_2 \geq \sigma_{k+1}$

Second: take $B = A_k$. Achieves the min. \square

Right and Left Singular vectors:

$$Av_i = \sigma_i u_i \\ A^T u_j = \sigma_j v_j$$

- Consequence $A^T Av_i = \sigma_i^2 v_i$ and $AA^T u_i = \sigma_i^2 u_i$
- Right singular vectors (v_i 's) are eigenvectors of $A^T A$
- Left singular vectors (u_i 's) are eigenvectors of AA^T
- Possible to get the SVD from eigenvectors of AA^T and $A^T A$ – but: difficulties due to non-uniqueness of the SVD

Define the $r \times r$ matrix

$$\Sigma_1 = \text{diag}(\sigma_1, \dots, \sigma_r)$$

- Let $A \in \mathbb{R}^{m \times n}$ and consider $A^T A$ ($\in \mathbb{R}^{n \times n}$):

$$A^T A = V \Sigma^T \Sigma V^T \rightarrow A^T A = V \underbrace{\begin{pmatrix} \Sigma_1^2 & 0 \\ 0 & 0 \end{pmatrix}}_{n \times n} V^T$$

- This gives the spectral decomposition of $A^T A$.

- Similarly, U gives the eigenvectors of AA^T .

$$AA^T = U \underbrace{\begin{pmatrix} \Sigma_1^2 & 0 \\ 0 & 0 \end{pmatrix}}_{m \times m} U^T$$

Important:

$A^T A = V D_1 V^T$ and $AA^T = U D_2 U^T$ give the SVD factors U, V up to signs!