

## Least-Squares Systems and The QR factorization

- Orthogonality
- Least-squares systems.
- The Gram-Schmidt and Modified Gram-Schmidt processes.
- The Householder QR and the Givens QR.

7-1

## Orthogonality

1. Two vectors  $u$  and  $v$  are orthogonal if  $(u, v) = 0$ .
  2. A system of vectors  $\{v_1, \dots, v_n\}$  is **orthogonal** if  $(v_i, v_j) = 0$  for  $i \neq j$ ; and **orthonormal** if  $(v_i, v_j) = \delta_{ij}$
  3. A matrix is **orthogonal** if its columns are orthonormal
- Notation:  $V = [v_1, \dots, v_n] \implies$  matrix with column-vectors  $v_1, \dots, v_n$ .
  - Orthogonality is essential in understanding and solving least-squares problems.

7-2

GvL 5, 5.3 – QR

7-2

## Least-Squares systems

- Given: an  $m \times n$  matrix  $n < m$ . Problem: find  $x$  which minimizes:

$$\|b - Ax\|_2$$

- Good illustration: Data fitting.

Typical problem of data fitting: We seek an unknown function as a linear combination  $\phi$  of  $n$  known functions  $\phi_i$  (e.g. polynomials, trig. functions). Experimental data (not accurate) provides measures  $\beta_1, \dots, \beta_m$  of this unknown function at points  $t_1, \dots, t_m$ . Problem: find the 'best' possible approximation  $\phi$  to this data.

$$\phi(t) = \sum_{i=1}^n \xi_i \phi_i(t) \quad , \quad \text{s.t.} \quad \phi(t_j) \approx \beta_j, \quad j = 1, \dots, m$$

7-3

GvL 5, 5.3 – QR

7-3

- Question: Close in what sense?
- Least-squares approximation: Find  $\phi$  such that

$$\phi(t) = \sum_{i=1}^n \xi_i \phi_i(t), \quad \& \quad \sum_{j=1}^m |\phi(t_j) - \beta_j|^2 = \text{Min}$$

- In linear algebra terms: find 'best' approximation to a vector  $b$  from linear combinations of vectors  $f_i, i = 1, \dots, n$ , where

$$b = \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_m \end{pmatrix}, \quad f_i = \begin{pmatrix} \phi_i(t_1) \\ \phi_i(t_2) \\ \vdots \\ \phi_i(t_m) \end{pmatrix}$$

7-4

GvL 5, 5.3 – QR

7-4

► We want to find  $x = \{\xi_i\}_{i=1,\dots,n}$  such that


$$\left\| \sum_{i=1}^n \xi_i f_i - b \right\|_2 \quad \text{Minimum}$$

Define

$$F = [f_1, f_2, \dots, f_n], \quad x = \begin{pmatrix} \xi_1 \\ \vdots \\ \xi_n \end{pmatrix}$$

► We want to find  $x$  to **minimize  $\|b - Fx\|_2$**

► This is a **Least-squares linear system**:  $F$  is  $m \times n$ , with  $m \geq n$ .

 Formulate the least-squares system for the problem of finding the polynomial of degree 2 that approximates a function  $f$  which satisfies  $f(-1) = -1; f(0) = 1; f(1) = 2; f(2) = 0$

**Solution:**  $\phi_1(t) = 1; \phi_2(t) = t; \phi_3(t) = t^2;$

• Evaluate the  $\phi_i$ 's at points  $t_1 = -1; t_2 = 0; t_3 = 1; t_4 = 2$ :

$$f_1 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \quad f_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \\ 2 \end{pmatrix} \quad f_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 4 \end{pmatrix} \quad \rightarrow$$

► So the coefficients  $\xi_1, \xi_2, \xi_3$  of the polynomial  $\xi_1 + \xi_2 t + \xi_3 t^2$  are the solution of the least-squares problem  $\min \|b - Fx\|$  where:

$$F = \begin{pmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{pmatrix} \quad b = \begin{pmatrix} -1 \\ 1 \\ 2 \\ 0 \end{pmatrix}$$


**THEOREM.** The vector  $x_*$  minimizes  $\psi(x) = \|b - Fx\|_2^2$  if and only if it is the solution of the **normal equations**:

$$F^T F x = F^T b$$

*Proof:* Expand out the formula for  $\psi(x_* + \delta x)$ :

$$\begin{aligned} \psi(x_* + \delta x) &= ((b - Fx_*) - F\delta x)^T ((b - Fx_*) - F\delta x) \\ &= \psi(x_*) - 2(F\delta x)^T (b - Fx_*) + (F\delta x)^T (F\delta x) \\ &= \psi(x_*) - 2(\delta x)^T \underbrace{[F^T (b - Fx_*)]}_{-\nabla_x \psi} + \underbrace{(F\delta x)^T (F\delta x)}_{\text{always } \geq 0} \end{aligned}$$


Can see that  $\psi(x_* + \delta x) \geq \psi(x_*)$  for any  $\delta x$ , iff the boxed quantity [the gradient vector] is zero. Q.E.D.



**Illustration of theorem:**  $x^*$  is the best approximation to the vector  $b$  from the subspace  $\text{span}\{F\}$  if and only if  $b - Fx^*$  is  $\perp$  to the whole subspace  $\text{span}\{F\}$ . This in turn is equivalent to  $F^T (b - Fx^*) = 0$  ► Normal equations.

**Example:**

Points:	$t_1 = -1$	$t_2 = -1/2$	$t_3 = 0$	$t_4 = 1/2$	$t_5 = 1$
Values:	$\beta_1 = 0.1$	$\beta_2 = 0.3$	$\beta_3 = 0.3$	$\beta_4 = 0.2$	$\beta_5 = 0.0$



1) Approximations by polynomials of degree one:

➤  $\phi_1(t) = 1, \phi_2(t) = t$ .


$$F = \begin{pmatrix} 1.0 & -1.0 \\ 1.0 & -0.5 \\ 1.0 & 0 \\ 1.0 & 0.5 \\ 1.0 & 1.0 \end{pmatrix}$$

$$F^T F = \begin{pmatrix} 5.0 & 0 \\ 0 & 2.5 \end{pmatrix}$$

$$F^T b = \begin{pmatrix} 0.9 \\ -0.15 \end{pmatrix}$$

➤ Best approximation is

$$\phi(t) = 0.18 - 0.06t.$$




2) Approximation by polynomials of degree 2:

➤  $\phi_1(t) = 1, \phi_2(t) = t, \phi_3(t) = t^2$ .

➤ Best polynomial found:

$$0.3085714285 - 0.06 \times t - 0.2571428571 \times t^2$$



*Problem with Normal Equations*

➤ Condition number is high: if  $A$  is square and non-singular, then

$$\kappa_2(A) = \|A\|_2 \cdot \|A^{-1}\|_2 = \sigma_{\max}/\sigma_{\min}$$

$$\kappa_2(A^T A) = \|A^T A\|_2 \cdot \|(A^T A)^{-1}\|_2 = (\sigma_{\max}/\sigma_{\min})^2$$

➤ Example: Let  $A = \begin{pmatrix} 1 & 1 & -\epsilon \\ \epsilon & 0 & 1 \\ 0 & \epsilon & 1 \end{pmatrix}$ .

➤ Then  $\kappa(A) = \sqrt{2}/\epsilon$ , but  $\kappa(A^T A) = 2\epsilon^{-2}$ .

➤  $fl(A^T A) = fl \begin{pmatrix} 1 + \epsilon^2 & 1 & 0 \\ 1 & 1 + \epsilon^2 & 0 \\ 0 & 0 & 2 + \epsilon^2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$

is singular to working precision (if  $\epsilon < \underline{u}$ ).

## Finding an orthonormal basis of a subspace

- Goal: Find vector in  $\text{span}(X)$  closest to  $b$ .
- Much easier with an orthonormal basis for  $\text{span}(X)$ .

**Problem:** Given  $X = [x_1, \dots, x_n]$ , compute  $Q = [q_1, \dots, q_n]$  which has orthonormal columns and s.t.  $\text{span}(Q) = \text{span}(X)$

- Note: each column of  $X$  must be a linear combination of certain columns of  $Q$ .
- We will find  $Q$  so that  $x_j$  ( $j$  column of  $X$ ) is a linear combination of the first  $j$  columns of  $Q$ .

7-13

GvL 5, 5.3 – QR

7-13

## ALGORITHM : 1. Classical Gram-Schmidt

1. For  $j = 1, \dots, n$  Do:
2. Set  $\hat{q} := x_j$
3. Compute  $r_{ij} := (\hat{q}, q_i)$ , for  $i = 1, \dots, j - 1$
4. For  $i = 1, \dots, j - 1$  Do :
5. Compute  $\hat{q} := \hat{q} - r_{ij}q_i$
6. EndDo
7. Compute  $r_{jj} := \|\hat{q}\|_2$ ,
8. If  $r_{jj} = 0$  then Stop, else  $q_j := \hat{q}/r_{jj}$
9. EndDo

- All  $n$  steps can be completed iff  $x_1, x_2, \dots, x_n$  are linearly independent.

 Prove this result

7-14

GvL 5, 5.3 – QR

7-14

- Lines 5 and 7-8 show that

$$x_j = r_{1j}q_1 + r_{2j}q_2 + \dots + r_{jj}q_j$$

- If  $X = [x_1, x_2, \dots, x_n]$ ,  $Q = [q_1, q_2, \dots, q_n]$ , and if  $R$  is the  $n \times n$  upper triangular matrix

$$R = \{r_{ij}\}_{i,j=1,\dots,n}$$

then the above relation can be written as

$$X = QR$$

- $R$  is upper triangular,  $Q$  is orthogonal. This is called the *QR factorization* of  $X$ .

 What is the cost of the factorization when  $X \in \mathbb{R}^{m \times n}$ ?

7-15

GvL 5, 5.3 – QR

7-15

$$\begin{array}{c}
 \boxed{X} \\
 \text{Original} \\
 \text{matrix}
 \end{array}
 =
 \begin{array}{c}
 \boxed{Q} \\
 \text{Q is orthogonal} \\
 (Q^H Q = I)
 \end{array}
 *
 \begin{array}{c}
 \boxed{R} \\
 \text{R is upper} \\
 \text{triangular}
 \end{array}$$

Another decomposition:

A matrix  $X$ , with linearly independent columns, is the product of an orthogonal matrix  $Q$  and a upper triangular matrix  $R$ .

7-16

GvL 5, 5.3 – QR

7-16

- Better algorithm: Modified Gram-Schmidt.

### ALGORITHM : 2. Modified Gram-Schmidt

1. For  $j = 1, \dots, n$  Do:
2.   Define  $\hat{q} := x_j$
3.   For  $i = 1, \dots, j - 1$ , Do:
4.      $r_{ij} := (\hat{q}, q_i)$
5.      $\hat{q} := \hat{q} - r_{ij}q_i$
6.   EndDo
7.   Compute  $r_{jj} := \|\hat{q}\|_2$ ,
8.   If  $r_{jj} = 0$  then Stop, else  $q_j := \hat{q}/r_{jj}$
9. EndDo

Only difference: inner product uses the accumulated subsum instead of original  $\hat{q}$

7-17


GvL 5, 5.3 – QR

7-17

- The operations in lines 4 and 5 can be written as

$$\hat{q} := ORTH(\hat{q}, q_i)$$

where  $ORTH(x, q)$  denotes the operation of orthogonalizing a vector  $x$  against a unit vector  $q$ .



Result of  $z = ORTH(x, q)$

7-18

GvL 5, 5.3 – QR

7-18

- Modified Gram-Schmidt algorithm is much more stable than classical Gram-Schmidt in general.

Suppose MGS is applied to  $A$  yielding computed matrices  $\hat{Q}$  and  $\hat{R}$ . Then there are constants  $c_i$  (depending on  $(m, n)$ ) such that

$$A + E_1 = \hat{Q}\hat{R} \quad \|E_1\|_2 \leq c_1 \underline{u} \|A\|_2$$

$$\|\hat{Q}^T \hat{Q} - I\|_2 \leq c_2 \underline{u} \kappa_2(A) + O((\underline{u} \kappa_2(A))^2)$$

for a certain perturbation matrix  $E_1$ , and there exists an orthonormal matrix  $Q$  such that

$$A + E_2 = Q\hat{R} \quad \|E_2(:, j)\|_2 \leq c_3 \underline{u} \|A(:, j)\|_2$$

for a certain perturbation matrix  $E_2$ .

7-19

GvL 5, 5.3 – QR

7-19

- An equivalent version:

### ALGORITHM : 3. Modified Gram-Schmidt - 2 -

0. Set  $\hat{Q} := X$
1. For  $i = 1, \dots, n$  Do:
2.   Compute  $r_{ii} := \|\hat{q}_i\|_2$ ,
3.   If  $r_{ii} = 0$  then Stop, else  $q_i := \hat{q}_i/r_{ii}$
4.   For  $j = i + 1, \dots, n$ , Do:
5.      $r_{ij} := (\hat{q}_j, q_i)$
6.      $\hat{q}_j := \hat{q}_j - r_{ij}q_i$
7.   EndDo
8. EndDo

- Does exactly the same computation as previous algorithm, but in a different order.

7-20

GvL 5, 5.3 – QR

7-20

**Example:**

Orthonormalize the system of vectors:

$$X = [x_1, x_2, x_3] = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & -1 \\ 1 & 0 & 4 \end{pmatrix}$$

Answer:

$$q_1 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}; \quad \hat{q}_2 = x_2 - (x_2, q_1)q_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} - 1 \times \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}$$

$$\hat{q}_2 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}; \quad q_2 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}$$

$$\hat{q}_3 = x_3 - (x_3, q_1)q_1 = \begin{pmatrix} 1 \\ 0 \\ -1 \\ 4 \end{pmatrix} - 2 \times \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \\ -2 \\ 3 \end{pmatrix}$$

$$\hat{q}_3 = \hat{q}_3 - (\hat{q}_3, q_2)q_2 = \begin{pmatrix} 0 \\ -1 \\ -2 \\ 3 \end{pmatrix} - (-1) \times \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \\ -\frac{1}{2} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \\ -2.5 \\ 2.5 \end{pmatrix}$$

$$\|\hat{q}_3\|_2 = \sqrt{13} \rightarrow q_3 = \frac{\hat{q}_3}{\|\hat{q}_3\|_2} = \frac{1}{\sqrt{13}} \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \\ -2.5 \\ 2.5 \end{pmatrix}$$

 For this example: what is  $Q$ ? what is  $R$ ? Compute  $Q^T Q$ .

➤ Result is the identity matrix.

Recall: For any orthogonal matrix  $Q$ , we have

$$Q^T Q = I$$

(In complex case:  $Q^H Q = I$ ).

Consequence: For an  $n \times n$  orthogonal matrix  $Q^{-1} = Q^T$ .  
( $Q$  is orthogonal/ unitary)

*Use of the QR factorization*

Problem:  $Ax \approx b$  in least-squares sense

$A$  is an  $m \times n$  (full-rank) matrix. Let

$$A = QR$$

the QR factorization of  $A$  and consider the normal equations:

$$A^T Ax = A^T b \rightarrow R^T Q^T QRx = R^T Q^T b \rightarrow$$

$$R^T Rx = R^T Q^T b \rightarrow Rx = Q^T b$$

( $R^T$  is an  $n \times n$  nonsingular matrix). Therefore,

$$x = R^{-1} Q^T b$$

**Another derivation:**

- Recall:  $\text{span}(Q) = \text{span}(A)$
- So  $\|b - Ax\|_2$  is minimum when  $b - Ax \perp \text{span}\{Q\}$
- Therefore solution  $x$  must satisfy  $Q^T(b - Ax) = 0 \rightarrow$   
 $Q^T(b - QRx) = 0 \rightarrow Rx = Q^Tb$

$$x = R^{-1}Q^Tb$$

- Also observe that for any vector  $w$

$$w = QQ^T w + (I - QQ^T)w$$

and that  $QQ^T w \perp (I - QQ^T)w \rightarrow$

- Pythagoras theorem  $\rightarrow \|w\|_2^2 = \|QQ^T w\|_2^2 + \|(I - QQ^T)w\|_2^2$


$$\begin{aligned} \|b - Ax\|^2 &= \|b - QRx\|^2 \\ &= \|(I - QQ^T)b + Q(Q^Tb - Rx)\|^2 \\ &= \|(I - QQ^T)b\|^2 + \|Q(Q^Tb - Rx)\|^2 \\ &= \|(I - QQ^T)b\|^2 + \|Q^Tb - Rx\|^2 \end{aligned}$$


- Min is reached when 2nd term of r.h.s. is zero.

Method:

- Compute the QR factorization of  $A$ ,  $A = QR$ .
- Compute the right-hand side  $f = Q^Tb$
- Solve the upper triangular system  $Rx = f$ .
- $x$  is the least-squares solution

- As a rule it is not a good idea to form  $A^T A$  and solve the normal equations. Methods using the QR factorization are better.

 5 Total cost?? (depends on the algorithm used to get the QR decomposition).

 6 Using matlab find the parabola that fits the data in previous data fitting example (p. 7-9) in L.S. sense [verify that the result found is correct.]

**Application:** another method for solving linear systems.

$$Ax = b$$

$A$  is an  $n \times n$  nonsingular matrix. Compute its QR factorization.

- Multiply both sides by  $Q^T \rightarrow Q^TQRx = Q^Tb \rightarrow$

$$Rx = Q^Tb$$

Method:

- Compute the QR factorization of  $A$ ,  $A = QR$ .
- Solve the upper triangular system  $Rx = Q^Tb$ .

 7 Cost??